


L5 ANSWER 1 OF 1 REGISTRY COPYRIGHT 2003 ACS  
 RN 298-14-6 REGISTRY  
 CN Carbonic acid, monopotassium salt (8CI, 9CI) (CA INDEX NAME)  
 OTHER NAMES:  
 CN Armicarb  
 CN Hydrogen potassium carbonate  
 CN K-Lyte  
 CN Kafylox  
 CN Kaligreen  
 CN Monopotassium carbonate  
 CN Potassium acid carbonate  
 CN **Potassium bicarbonate**  
 CN Potassium bicarbonate (KHCO3)  
 CN Potassium carbonate (KHCO3)  
 CN Potassium hydrogen carbonate (KHCO3)  
 CN Purple K  
 MF C H2 O3 . K  
 CI COM  
 LC STN Files: ADISNEWS, AGRICOLA, ANABSTR, BEILSTEIN\*, BIOBUSINESS, BIOSIS,  
 BIOTECHNO, CA, CANCERLIT, CAOLD, CAPLUS, CASREACT, CBNB, CHEMCATS,  
 CHEMINFORMRX, CHEMLIST, CIN, CSCHM, DDFU, DETHERM\*, DIOGENES, DRUGU,  
 EMBASE, ENCOMPLIT, ENCOMPLIT2, ENCOMPPAT, ENCOMPPAT2, GMELIN\*, IFICDB,  
 IFIPAT, IFIUDB, IPA, MEDLINE, MRCK\*, MSDS-OHS, NIOSHTIC, PDLCOM\*, PHAR,  
 PIRA, PROMT, TOXCENTER, TULSA, USAN, USPAT2, USPATFULL, VTB  
 (\*File contains numerically searchable property data)  
 Other Sources: DSL\*\*, EINECS\*\*, TSCA\*\*  
 (\*\*Enter CHEMLIST File for up-to-date regulatory information)  
 CRN (463-79-6)



● K

3285 REFERENCES IN FILE CA (1957 TO DATE)  
 35 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA  
 3287 REFERENCES IN FILE CAPLUS (1957 TO DATE)  
 3 REFERENCES IN FILE CAOLD (PRIOR TO 1967)

=>

=> d

L6 ANSWER 1 OF 1 REGISTRY COPYRIGHT 2003 ACS

RN 71-52-3 REGISTRY

CN Carbonate, hydrogen (8CI, 9CI) (CA INDEX NAME)

OTHER NAMES:

CN Bicarbonate

CN Bicarbonate (HCO<sub>3</sub><sup>-</sup>)

CN Bicarbonate anion

CN **Bicarbonate ion**

CN Bicarbonate ion (HCO<sub>3</sub><sup>1-</sup>)

CN Carbonate (HCO<sub>3</sub><sup>1-</sup>)

CN Carbonate ion (HCO<sub>3</sub><sup>1-</sup>)

CN Carbonic acid, ion(1-)

CN Hydrocarbonate(1-)

CN Hydrogen carbonate

CN Hydrogen carbonate (HCO<sub>3</sub><sup>-</sup>)

CN Hydrogen carbonate anion

CN Hydrogen carbonate ion

CN Hydrogen carbonate ion (HCO<sub>3</sub><sup>-</sup>)


CN Monohydrogen carbonate

MF C H O<sub>3</sub>

CI COM

LC STN Files: ADISNEWS, AGRICOLA, ANABSTR, BEILSTEIN\*, BIOBUSINESS, BIOSIS, BIOTECHNO, CA, CAPLUS, CASREACT, CBNB, CEN, CHEMINFORMRX, CIN, EMBASE, GMELIN\*, IFICDB, IFIPAT, IFIUDB, NIOSHTIC, PIRA, PROMT, SPECINFO, TOXCENTER, TULSA, USPAT2, USPATFULL

(\*File contains numerically searchable property data)



12700 REFERENCES IN FILE CA (1957 TO DATE)

98 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA

12710 REFERENCES IN FILE CAPLUS (1957 TO DATE)

=>

L2 ANSWER 1 OF 1 REGISTRY COPYRIGHT 2003 ACS

RN 63-42-3 REGISTRY

CN D-Glucose, 4-O-.beta.-D-galactopyranosyl- (9CI) (CA INDEX NAME)

OTHER CA INDEX NAMES:

CN Lactose (8CI)

OTHER NAMES:

CN (+)-Lactose

CN AHL

CN Aletobiose

CN D-(+)-Lactose

CN Fast-flo

CN Fast-Flo Lactose

CN Galactinum

CN Lactin

CN Lactin (carbohydrate)

CN Lactobiose

CN Lactose anhydride

CN Lactose anhydrous

CN Lactose Fast-flo

CN Milk sugar

CN Nonpareil 107

CN Osmolactan

CN Pharmatose 21

CN Pharmatose 325M

CN Pharmatose 450M

CN Saccharum lacticum

CN Tablettose

CN Tablettose 70

CN Zeparox EP

AR 16984-38-6

FS STEREOSEARCH

DR 1336-90-9, 36570-80-6, 73824-63-2, 89466-76-2, 35396-14-6

MF C12 H22 O11

CI COM


LC STN Files: ADISNEWS, AGRICOLA, ANABSTR, BEILSTEIN\*, BIOBUSINESS, BIOSIS, BIOTECHNO, CA, CANCERLIT, CAOLD, CAPLUS, CASREACT, CBNB, CEN, CHEMCATS, CHEMLIST, CHEMSAFE, CIN, CSCHM, CSNB, DDFU, DETHERM\*, DIOGENES, DRUGU, EMBASE, IFICDB, IFIPAT, IFIUDB, IPA, MEDLINE, MRCK\*, MSDS-OHS, NIOSHTIC, PDLCOM\*, PIRA, PROMT, RTECS\*, SPECINFO, TOXCENTER, TULSA, USPAT2, USPATFULL, VETU

(\*File contains numerically searchable property data)

Other Sources: DSL\*\*, EINECS\*\*, TSCA\*\*

(\*\*Enter CHEMLIST File for up-to-date regulatory information)

Absolute stereochemistry. Rotation (+).




\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

> d 1 2

L1 ANSWER 1 OF 2 REGISTRY COPYRIGHT 2003 ACS  
RN 87-78-5 REGISTRY  
CN Mannitol (8CI, 9CI) (CA INDEX NAME)  
OTHER NAMES:  
CN Mannidex 16700  
FS STEREOSEARCH  
DR 133-43-7, 36413-61-3, 5149-40-6  
MF C6 H14 O6  
CI COM  
LC STN Files: ADISNEWS, AGRICOLA, ANABSTR, AQUIRE, BEILSTEIN\*, BIOBUSINESS,  
BIOSIS, BIOTECHNO, CA, CAOLD, CAPLUS, CASREACT, CBNB, CEN, CHEMCATS,  
CHEMINFORMRX, CHEMLIST, CIN, DETHERM\*, DIOGENES, EMBASE, GMELIN\*,  
HODOC\*, IFICDB, IFIPAT, IFIUDB, MEDLINE, NAPRALERT, NIOSHTIC, PDLCOM\*,  
PHARMASEARCH, PIRA, PROMT, RTECS\*, TOXCENTER, TULSA, USPAT2, USPATFULL  
(\*File contains numerically searchable property data)  
Other Sources: DSL\*\*, EINECS\*\*, TSCA\*\*  
(\*\*Enter CHEMLIST File for up-to-date regulatory information)

Relative stereochemistry.



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

165 REFERENCES IN FILE CA (1957 TO DATE)  
12 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA  
167 REFERENCES IN FILE CAPLUS (1957 TO DATE)  
1 REFERENCES IN FILE CAOLD (PRIOR TO 1967)

L1 ANSWER 2 OF 2 REGISTRY COPYRIGHT 2003 ACS  
RN 69-65-8 REGISTRY  
CN D-Mannitol (9CI) (CA INDEX NAME)  
OTHER CA INDEX NAMES:  
CN Cordycepic acid (6CI, 7CI)  
CN Mannitol, D- (8CI)  
OTHER NAMES:  
CN D-(-)-Mannitol  
CN Diosmol  
CN Isotol  
CN Manicol  
CN Maniton S  
CN Manna sugar  
CN Mannidex  
CN Mannigen  
CN Mannistol  
CN Mannit  
CN Mannite  
CN Mannitol  
CN Mannitolium  
CN Mannogem 2080  
CN Marine Crystal  
CN Osmitrol  
CN Osmosal  
CN Resectisol  
FS STEREOSEARCH

DR 123897-58-5, 75398-80-0, 85085-15-0

MF C6 H14 O6

CI COM


LC STN Files: ADISNEWS, AGRICOLA, ANABSTR, BEILSTEIN\*, BIOBUSINESS, BIOSIS, BIOTECHNO, CA, CABA, CANCERLIT, CAOLD, CAPLUS, CASREACT, CEN, CHEMCATS, CHEMINFORMRX, CHEMLIST, CIN, CSChem, CSNB, DDFU, DETHERM\*, DIOGENES, DRUGU, EMBASE, GMELIN\*, HODOC\*, HSDB\*, IFICDB, IFIPAT, IFIUDb, IPA, MEDLINE, MRCK\*, MSDS-OHS, NAPRALERT, NIOSHTIC, PDLCOM\*, PHARMASEARCH, PIRA, PROMT, RTECS\*, SPECINFO, TOXCENTER, TULSA, USAN, USPAT2, USPATFULL, VETU

(\*File contains numerically searchable property data)

Other Sources: DSL\*\*, EINECS\*\*, TSCA\*\*

(\*\*Enter CHEMLIST File for up-to-date regulatory information)

Absolute stereochemistry.



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

12451 REFERENCES IN FILE CA (1957 TO DATE)

283 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA

12479 REFERENCES IN FILE CAPLUS (1957 TO DATE)

2 REFERENCES IN FILE CAOLD (PRIOR TO 1967)

=>

L61 ANSWER 31 OF 294 CA COPYRIGHT 2003 ACS

AN 136:147485 CA

TI An advantageous carrier **solution** for vitrifiable concentrations of cryoprotectants, and compatible cryoprotectant mixtures

IN Fahy, Gregory M.; Wowk, Brian

PA 21st Century Medicine, USA

SO PCT Int. Appl., 16 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 4

PATENT NO. KIND DATE APPLICATION NO. DATE

PI WO 2002009516 A2 20020207 WO 2001-US23853 20010730

WO 2002009516 A3 20020613

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM

RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

EP 1311155 A2 20030521 EP 2001-967949 20010730

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR

PRAI US 2000-221691P P 20000731

WO 2001-US23853 W 20010730

AB Disclosed herein is a carrier **soln.** for cryoprotectants that is useful for use with cells, tissues, and whole organs and for a variety of cryoprotectant **solns.** and that permits antinucleators to be fully effective in vitrification **solns.**, thereby allowing vitrification **solns.** to attain extreme effectiveness, and compatible vitrification **soln.** compns. for use with this carrier **soln.** The carrier **soln.** comprises lactose and mannitol as well as other beneficial ingredients.

IC ICM A01N001-02

CC 9-11 (Biochemical Methods)

ST carrier **soln** vitrifiable concn cryoprotectant compatible

IT Animal tissue

Carriers

Cell

Concentration (condition)

Cryopreservation

Cryoprotectants

Kidney

Mixtures

Organ, animal

**Solutions**

Vitrification


Washing

(advantageous carrier **soln.** for vitrifiable concns. of cryoprotectants, and compatible cryoprotectant mixts.)

IT 50-99-7, Glucose, biological studies 57-50-1, Sucrose, biological studies 63-42-3, Lactose 67-68-5, Dimethyl sulfoxide, biological studies 69-65-8, Mannitol 75-12-7, Formamide, biological studies 107-21-1, Ethylene glycol, biological studies 9002-89-5, Polyvinyl alcohol 9003-20-7D, Poly(vinyl acetate), 80% hydrolyzed 9003-39-8, Polyvinylpyrrolidone 9041-07-0, Decaglycerol 25213-24-5 25618-55-7, Polyglycerol 394248-23-8, X 1000

RL: BUU (Biological use, unclassified); BIOL (Biological study); USES

L3 ANSWER 1 OF 1 REGISTRY COPYRIGHT 2003 ACS  
 RN 144-55-8 REGISTRY  
 CN Carbonic acid monosodium salt (8CI, 9CI) (CA INDEX NAME)  
 OTHER NAMES:  
 CN Baking soda  
 CN BI-CF 40E  
 CN BI-H 40E  
 CN Carbonic acid sodium salt (1:1)  
 CN Carbonic acid, monosodium salt  
 CN Cellborn SC-K  
 CN Cellborn SC-P  
 CN Cellmic 266  
 CN DP 35/22  
 CN Extin B  
 CN Meylon  
 CN Monosodium carbonate  
 CN Monosodium hydrogen carbonate  
 CN Soda  
 CN Sodium acid carbonate  
 CN **Sodium bicarbonate**  
 CN Sodium carbonate (Na(HCO<sub>3</sub>))  
 CN Sodium hydrogen carbonate  
 CN Sodium monohydrogen carbonate  
 CN Soludal  
 CN Unifine P 4  
 DR 196216-68-9, 199723-76-7, 246180-97-2  
 MF C H2 O3 . Na  
 CI COM  
 LC STN Files: ADISNEWS, AGRICOLA, ANABSTR, AQUIRE, BEILSTEIN\*, BIOBUSINESS,  
 BIOSIS, BIOTECHNO, CA, CABA, CANCERLIT, CAOLD, CAPLUS, CASREACT, CBNB,  
 CEN, CHEMCATS, CHEMINFORMRX, CHEMLIST, CIN, CSCHM, CSNB, DDFU,  
 DETHERM\*, DIOGENES, DIPPR\*, DRUGU, EMBASE, ENCOMPLIT, ENCOMPLIT2,  
 ENCOMPPAT, ENCOMPPAT2, GMELIN\*, HSDB\*, IFICDB, IFIPAT, IFIUDB, IPA,  
 MEDLINE, MRCK\*, MSDS-OHS, NIOSHTIC, PDLCOM\*, PHARMASEARCH, PIRA, PROMT,  
 RTECS\*, TOXCENTER, TULSA, USAN, USPAT2, USPATFULL, VETU, VTB  
 (\*File contains numerically searchable property data)  
 Other Sources: DSL\*\*, EINECS\*\*, TSCA\*\*  
 (\*\*Enter CHEMLIST File for up-to-date regulatory information)  
 CRN (463-79-6)



● Na

*inverted 211 - 42 - 51*  
*factor 113 - 37 52*  
*bicarb 125 - 135 53*

methacrylate polymer 9003-53-6, Polystyrene 9003-54-7,  
Acrylonitrile-styrene copolymer 9003-56-9 9010-79-1,  
Ethylene-propylene copolymer 9011-14-7, Methyl methacrylate polymer  
24937-78-8, Ethylene-vinyl acetate copolymer 24968-12-5, Polybutylene  
terephthalate 25014-41-9, Acrylonitrile polymer 25038-59-9,  
Polyethylene terephthalate, biological studies 25067-61-2,  
Methacrylonitrile polymer 25085-46-5, Ethylene-vinyl acetate-vinyl  
chloride copolymer 26062-94-2, Polybutylene terephthalate 120460-98-2,  
Urethane-vinyl chloride copolymer

RL: BIOL (Biological study)  
(medical goods contg. hemolysis inhibitors and)

- L61 ANSWER 230 OF 294 CA COPYRIGHT 2003 ACS  
AN 102:181076 CA  
TI Synergistic depression of the freezing temperature in **solutions**  
of polyhydroxy compounds and antifreeze glycoproteins  
AU Kerr, William L.; Burcham, Timothy S.; Osuga, David T.; Yeh, Yin; Feeney,  
Robert E.; Caple, Gerald  
CS Dep. Food Sci. Technol., Univ. California, Davis, CA, 95616, USA  
SO Cryo-Letters (1985), 6(2), 107-14  
CODEN: CRLED9; ISSN: 0143-2044  
DT Journal  
LA English  
AB Mixts. of various polyhydroxy compds. with the low-mol.-wt. fractions of  
antifreeze glycoprotein from Antarctic fish blood showed a greater than  
additive lowering of the freezing temp. Mixts. of polyhydroxy compds.  
with higher-mol.-wt. antifreeze glycoproteins showed a much smaller  
synergistic effect on the freezing temp.  
CC 6-3 (General Biochemistry)  
Section cross-reference(s): 12  
ST polyhydroxy compd antifreeze glycoprotein synergism; freezing point  
depression polyhydroxy compd glycoprotein; fish glycoprotein polyhydroxy  
compd synergism  
IT Pagothenia borchgrevinki  
(antifreeze glycoproteins of, f.p. depression by, polyhydroxy compds.  
synergism with)  
IT Carbohydrates and Sugars, properties  
RL: PRP (Properties)  
(f.p. depression response to antifreeze glycoprotein synergism with)  
IT Glycoproteins  
RL: BIOL (Biological study)  
(AFGP-4, f.p. depression by, polyhydroxy compds. synergism with)  
IT Glycopeptides  
Glycoproteins  
RL: BIOL (Biological study)  
(antifreeze, f.p. depression by, polyhydroxy compds. synergism with)  
IT Hydroxy compounds  
RL: BIOL (Biological study)  
(poly-, f.p. depression response to antifreeze glycoprotein synergism  
with)  
IT 50-70-4, properties 50-99-7, properties 56-81-5, properties  
57-50-1, properties 59-23-4, properties 63-42-3  
69-65-8 69-79-4 87-89-8 87-99-0 488-81-3  
RL: PRP (Properties)  
(f.p. depression by antifreeze glycoprotein synergism with)
- L61 ANSWER 250 OF 294 CA COPYRIGHT 2003 ACS  
AN 98:8173 CA  
TI Water-soluble preparation for making an isotonic nitroglycerin  
**solution**  
IN Muench, Ulrich; Giesselmann, Ewald  
PA Sanol Schwarz-Monheim G.m.b.H., Fed. Rep. Ger.  
SO Ger. Offen., 16 pp.  
CODEN: GWXXBX



DT Patent  
LA German  
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	DE 3109783	A1	19821007	DE 1981-3109783	19810313
	DE 3109783	C2	19870402		
	FR 2501675	A1	19820917	FR 1982-4091	19820311
	FR 2501675	B1	19860919		
	WO 8203172	A1	19820930	WO 1982-DE54	19820312
	W: AT, CH, GB, HU, JP, LU, NL, US				
	JP 58500327	T2	19830303	JP 1982-500862	19820312
	JP 06015470	B4	19940302		
	GB 2110535	A1	19830622	GB 1982-32543	19820312
	GB 2110535	B2	19850206		
	US 4481220	A	19841106	US 1982-438887	19820929
	AT 8802240	A	19890315	AT 1988-2240	19880913
	AT 389050	B	19891010		
PRAI	DE 1981-3109783		19810313		
	AT 1982-9015		19820312		
	WO 1982-DE54		19820312		
AB	A prepn. that can be dild. with H2O to give an isotonic infusion or injection soln. contg. 1 mg/mL nitroglycerin [55-63-0] contains nitroglycerin and a solubilizer in a ratio of 0.5:1 to 2:1 and a solid nitroglycerin carrier capable of controlling isotonicity. Thus, 1.6 kg nitroglycerin was dissolved in 8-10 L Et2O and mixed with 1.6 kg 1,2-propylene glycol [57-55-6]. The soln. was mixed with 76.8 kg glucose [50-99-7] with aeration until the Et2O evapd. and the soln. was homogeneous. The soln., 11.550 kg, was added gradually to H2O at 70.degree. until dissolved, cooled, and dild. with H2O (total of 220 L). The soln. was placed in ampuls or bottles for injection.				
IC	A61K031-21				
CC	63-6 (Pharmaceuticals)				
ST	nitroglycerin solubilizer injection; propylene glycol nitroglycerin injection				
IT	Solubilizers				
	(for nitroglycerin injections)				
IT	55-63-0				
	RL: BIOL (Biological study)				
	(injections, solubilizers and carriers for)				
IT	50-70-4, biological studies 50-99-7, biological studies 57-48-7, biological studies 63-42-3 69-65-8 7647-14-5, biological studies				
	RL: BIOL (Biological study)				
	(nitroglycerin injections contg., for controlling isotonicity)				
IT	51-79-6	97-64-3	100-79-8	107-88-0	111-55-7 111-96-6 112-60-7
	123-80-8	126-33-0	127-19-5	542-59-6	1187-03-7 1569-01-3
	4128-76-1	5422-34-4	5464-28-8	9003-11-6	9004-32-4 11111-34-5
	19354-27-9	24567-27-9	25322-68-3	29387-84-6	53778-73-7
	83931-54-8				
	RL: BIOL (Biological study)				
	(nitroglycerin solubilization by, for injections)				
IT	50-21-5, properties	56-81-5, properties	57-13-6, biological studies		
	57-55-6, properties	68-12-2, properties			
	RL: PRP (Properties)				
	(nitroglycerin solubilization by, for injections)				
L61	ANSWER 270 OF 294 CA COPYRIGHT 2003 ACS				
AN	85:18011 CA				
TI	Lyophilization of hemoglobin solutions. Study of protector compounds capable of preventing the formation of methemoglobin				
AU	Labrude, P.; Vigneron, C.; Streiff, F.				
CS	Cent. Reg. Transfus. Sang. Nancy-Brabois, Vandoeuvre, Fr.				

SO Journal de Pharmacie de Belgique (1976), 31(2), 191-8  
CODEN: JPBEAJ; ISSN: 0047-2166

DT Journal

LA French

AB The protection of Hb **solns.** against oxidn. to methemoglobin during lyophilization by plasma expanders, macromols., sugars, glycerol, and THAM was studied. The most active mols. were the sugars and THAM which almost entirely prevented the formation of methemoglobin at a concn. of 1.25%.

CC 13-5 (Mammalian Biochemistry)

ST Hb oxidn lyophilization preservative; methemoglobin formation lyophilization preservative

IT Blood substitutes  
Albumins  
Sugars, biological studies  
RL: BIOL (Biological study)  
(Hb oxidn. response to, in lyophilization)

IT Methemoglobins  
RL: FORM (Formation, nonpreparative)  
(formation of, in lyophilization, prevention of)

IT Freeze drying  
(of Hb, oxidn. prevention in)

IT Hemoglobins  
RL: RCT (Reactant); RACT (Reactant or reagent)  
(oxidn. of, in lyophilization, prevention of)

IT 50-70-4 50-99-7, biological studies 56-81-5, biological studies 57-48-7, biological studies 57-50-1, biological studies 59-23-4, biological studies 63-42-3 69-65-8 77-86-1 3458-28-4 8057-73-6 9002-89-5 9003-39-8 9004-54-0, biological studies 39290-10-3 54847-63-1 66455-30-9  
RL: BIOL (Biological study)  
(Hb oxidn. response to, in lyophilization)

L61 ANSWER 272 OF 294 CA COPYRIGHT 2003 ACS

AN 83:197730 CA

TI Comparative testing of titrable acidity degree and pH value of infusion **solutions**

AU Horvath, Klara; Regos, Erika; Varga, Sarolta

CS Inst. Serobacteriol. Prod. Res. "HUMAN", Godollo, Hung.

SO Annales Immunologiae Hungaricae (1973), 17, 265-8  
CODEN: AIMHA3; ISSN: 0570-1708

DT Journal

LA English

AB The titrable acidity degree (i.e. the hydrogen ion reserve possessed by the nondissociated species of weak acids) of various infusion **solns.** was reported and compared with the pH of the same **solns.** Since in the case of weak acids pH does not characterize unequivocally the acidic character of the **soln.** it was suggested that the labels of infusion **solns.** also contain the value of titrable acidity.

CC 63-5 (Pharmaceuticals)

ST infusion **soln** titrable acidity

IT Ringer's **solution**  
RL: BIOL (Biological study)  
(acetate infusion **soln.**, titrable acidity degree of, pH in relation to)

IT Pharmaceuticals  
(infusion **solns.**, titrable acidity degree of, pH in relation to)

IT Ringer's **solution**  
RL: BIOL (Biological study)  
(lactate infusion **soln.**, titrable acidity degree of, pH in relation to)

IT Ringer's **solution**

RL: BIOL (Biological study)  
(titrable acidity degree of, pH in relation to)

IT 9004-54-0, biological studies

RL: BIOL (Biological study)  
(infusion glucose soln., titrable acidity degree of, pH in relation to)

IT 50-99-7, biological studies 57-48-7, biological studies

63-42-3 69-65-8 9004-54-0, biological studies

57455-70-6 57455-71-7

RL: BIOL (Biological study)  
(infusion soln., titrable acidity degree of, pH in relation to)

=>

(Uses)

(advantageous carrier soln. for vitrifiable concns. of  
cryoprotectants, and compatible cryoprotectant mixts.)

L61 ANSWER 65 OF 294 CA COPYRIGHT 2003 ACS  
AN 133:109998 CA  
TI Beta-interferon lozenge and its preparing method  
IN Cao, Xuetao; Ju, Dianwen; Tao, Qun  
PA Huachen Biological Technology Inst., Shanghai, Peop. Rep. China  
SO Faming Zhuanli Shenqing Gongkai Shuomingshu, 13 pp.  
CODEN: CNXXEV  
DT Patent  
LA Chinese  
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	CN 1227124	A	19990901	CN 1998-105383	19980225
PRAI	CN 1998-105383		19980225		
AB	The beta interferon lozenge is composed of 1-10 kIU beta-interferon and medicinal adjuvant. The medicinal adjuvant is selected from <u>one or more</u> of human serum albumin, bovine serum protein, polyethylene glycol, mannitol, lactose, glucose, starch, Mg stearate, and dextrin. The beta-interferon lozenge may contain 100-1,000 IU alpha-interferon and/or 1-10 kIU interleukin-2. The beta interferon lozenge is prepd. by mixing medicinal adjuvant, sieving, drying to obtain blank granule; spraying beta-interferon soln. in the blank granule; drying, and tableting. The lozenge is used for treatment of virus infection and/or tumor.				
IC	ICM A61K038-21				
	ICS A61K009-20				
CC	63-6 (Pharmaceuticals)				
	Section cross-reference(s): 15				
ST	beta interferon lozenge prepn				
IT	Proteins, general, biological studies				
	RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)				
	(blood; beta-interferon lozenge and its prepg. method)				
IT	Drug delivery systems				
	(lozenges; beta-interferon lozenge and its prepg. method)				
IT	Interferons				
	RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)				
	(.alpha.; .beta.-interferon lozenge and its prepg. method)				
IT	Antitumor agents				
	Antiviral agents				
	(.beta.-interferon lozenge and its prepg. method)				
IT	Albumins, biological studies				
	Interleukin 2				
	Polyoxyalkylenes, biological studies				
	RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)				
	(.beta.-interferon lozenge and its prepg. method)				
IT	Interferons				
	RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)				
	(.beta.; .beta.-interferon lozenge and its prepg. method)				
IT	9004-53-9, Dextrin		9005-25-8, Starch, biological studies		
	RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)				
	(beta-interferon/lozenge and its prepg. method)				
IT	50-99-7, Glucose, biological studies		63-42-3, Lactose		
	69-65-8, Mannitol		557-04-0, Magnesium stearate		25322-68-3,
	Polyethylene glycol				
	RL: THU (Therapeutic use); BIOL (Biological study); USES (Uses)				
	(.beta.-interferon lozenge and its prepg. method)				

L61 ANSWER 191 OF 294 CA COPYRIGHT 2003 ACS  
AN 112:62604 CA  
TI Monocarboxylic acid esters as blood preservatives

IN Nagai, Hiroshi; Kubota, Yoshiki; Tamura, Yoko; Kimura, Akio  
PA Terumo Corp., Japan; Kao Corp.  
SO Jpn. Kokai Tokkyo Koho, 8 pp.  
CODEN: JKXXAF

DT Patent  
LA Japanese

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 01106826	A2	19890424	JP 1987-263695	19871021
PRAI	JP 1987-263695		19871021		

OS MARPAT 112:62604

AB A preservative for blood contains an antihemolytic agent, RCO2R1 (R and R1 = C .gtoreq.3 hydrocarbyl; the sum of C in R and R1 is 11-30), together with other preservatives such as Na citrate, citric acid, glucose, NaH2PO4, adenine, NaCl, mannitol, etc. Thus, Tween-80 was dissolved (600 .mu.g/mL) in a soln. contg. NaCl 140, adenine 1.25, and glucose 50 mM, and 12 mM iso-Pr isolaurate was added. This emulsion (1.0 mL) was added as a preservative to 2.0 mL human erythrocyte conc. (with hematocrit value 70%) and stored for 5 wk at 4.degree..

IC ICM A61K035-14

CC 63-3 (Pharmaceuticals)

ST blood preservative antihemolytic carboxylate

IT Blood preservatives

(contg. monocarboxylic acid esters, as antihemolytic agents)

IT Carboxylic acids, esters

RL: BIOL (Biological study)

(esters, blood preservatives contg., as antihemolytic agents)

IT 50-70-4, Sorbitol, biological studies 50-99-7, D-Glucose, biological studies 57-50-1, Sucrose, biological studies 63-42-3

Lactose 68-04-2, Sodium citrate 69-65-8, Mannitol 69-79-4,

Maltose 73-24-5, Adenine, biological studies 77-92-9, Citric acid, biological studies 585-88-6, Maltitol 7558-80-7, Monosodium phosphate 7647-14-5, Sodium chloride, biological studies

RL: BIOL (Biological study)

(blood preservatives contg. antihemolytic carboxylic esters and)

IT 112-11-8, Isopropyl oleate 7425-14-1, 2-Ethylhexyl 2-ethylhexanoate 81897-25-8, 2-Ethylhexyl isostearate 120470-79-3, Isopropyl isolaurate

RL: BIOL (Biological study)

(blood preservatives/contg., as antihemolytic agent)

L61 ANSWER 197 OF 294 CA COPYRIGHT 2003 ACS

AN 110:219123 CA

TI Hemolysis inhibitors and medical resin compositions, medical goods, and blood-preserving fluids containing the same

IN Nagai, Hirofumi; Kubota, Yoshinori; Tamura, Yoko; Sado, Mineo; Kora, Shinichi; Kimura, Akio; Suzue, Shigetoshi; Miyamoto, Norioki

PA Kao Corp., Japan; Terumo Corp.

SO PCT Int. Appl., 163 pp.

CODEN: PIXXD2

DT Patent

LA Japanese

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 8803027	A1	19880505	WO 1987-JP810	19871022
	W: AU, US				
	RW: BE, DE, FR, GB, IT, NL, SE				
	JP 63104916	A2	19880510	JP 1986-249552	19861022
	JP 04015203	B4	19920317		
	AU 8781050	A1	19880525	AU 1987-81050	19871022
	AU 623071	B2	19920507		
	ES 2005457	A6	19890301	ES 1987-3293	19871022
	EP 334956	A1	19891004	EP 1987-906934	19871022

R: BE, DE, FR, GB, IT, NL, SE

PRAI JP 1986-249552 19861022

WO 1987-JP810 19871022

- AB Hemolysis inhibitors contg. carboxylic acid ester polymers, ether compds., and/or monocarboxylic ester are used to manuf. clin. resin compns., medical goods and blood preservers. Iso-Pr isotridecyl maleate (I) was dispersed in a mixt. of polyoxyethylene monooleate-saline (2000 .mu.g/mL) to make a 4 mM emulsion. This emulsion was added to a fluid contg. human erythrocytes having 70% hematocrit value such that the concn. of I was 0.4 mM. The fluid kept at 4.degree. for 4 wk showed the stability of erythrocytes. The stability of erythrocytes was also demonstrated by keeping blood in a poly(vinyl chloride) bag which consisted of poly(vinyl chloride) 100, didecyl phthalate 30, iso-Pr decyl maleate 20, epoxylated soybean oil 10, and a Ca-Zn type stabilizer 0.1 parts by wt.
- IC ICM A61K035-14  
ICS A61L031-00; A61M005-00; A01N001-02
- CC 63-7 (Pharmaceuticals)
- ST hemolysis inhibitor medical goods; blood preservative hemolysis inhibitor
- IT Glycerides, biological studies  
RL: DEV (Device component use); USES (Uses)  
(as hemolysis inhibitors, medical goods contg.)
- IT Blood preservation  
(carboxylic acid esters and ethers in, as hemolysis inhibitors)
- IT Medical goods  
(hemolysis inhibitor-contg.)
- IT Hemolysis  
(inhibitors, carboxylic acid derivs. as)
- IT Acrylic polymers, biological studies  
Polycarbonates, biological studies  
Polyesters, biological studies  
Urethane polymers, biological studies  
RL: BIOL (Biological study)  
(medical goods contg. hemolysis inhibitors and)
- IT Erythrocyte  
(stabilization of, by carboxylic acid derivs.)
- IT 50-70-4, D-Glucitol, biological studies 50-99-7, D-Glucose, biological studies 57-50-1, biological studies 63-42-3, Lactose 68-04-2 69-65-8, Mannitol 69-79-4 73-24-5, Adenine, biological studies 77-92-9, Citric acid, biological studies 585-88-6, Maltitol 7558-80-7, Monosodium phosphate 7647-14-5, Sodium chloride, biological studies  
RL: BIOL (Biological study)  
(as blood preservative soln. contg.)
- IT 120486-11-5  
RL: BIOL (Biological study)  
(as hemolysis inhibitor)
- IT 112-11-8, Isopropyl oleate 7425-14-1, 2-Ethylhexyl 2-ethylhexanoate 59068-03-0, Glycerin 1,3-bis(2-ethylhexyl) ether 81897-25-8 93120-93-5, Glycerin 1-butyl-3-isostearyl ether 120422-95-9 120422-96-0 120470-78-2 120470-79-3 120573-91-3  
RL: DEV (Device component use); USES (Uses)  
(as hemolysis inhibitor, medical goods contg.)
- IT 97-65-4D, esters 110-15-6D, Butanedioic acid, esters 110-16-7D, Maleic acid, esters 110-94-1D, Glutaric acid, esters 111-20-6D, Sebacic acid, esters 124-04-9D, Hexanedioic acid, esters 143-07-7D, Dodecanoic acid, esters 328-42-7D, Oxalacetic acid, esters  
RL: DEV (Device component use); USES (Uses)  
(as hemolysis inhibitors, medical goods contg.)
- IT 84-77-5, Didecyl phthalate 89-04-3, Trioctyl trimellitate  
RL: BIOL (Biological study)  
(as plasticizer, medical goods resin compns. contg.)
- IT 9002-86-2, Poly(vinyl chloride) 9002-88-4, Polyethylene 9003-07-0, Polypropylene 9003-21-8, Methyl acrylate polymer 9003-22-9, Vinyl acetate-vinyl chloride copolymer 9003-32-1 9003-42-3, Ethyl

L69 ANSWER 24 OF 39 WPIDS (C) 2003 THOMSON DERWENT  
AN 1998-076784 [07] WPIDS  
CR 1998-593980 [50]  
DNC C1998-025629  
TI **Solution** for preservation of biological materials - comprises two neutral solutes, especially raffinose and tri methylamine oxide. 103  
DC B04 D16 D22 E19 E37 P31  
IN FERGUSON, A B; WIGGINS, P M; WATSON, J D  
PA (BIOS-N) BIOSTORE NEW ZEALAND LTD; (BIOS-N) BIOSTORE NZ LTD; (BIOS-N) BIOSTORE NEW ZEALAND 0.5  
CYC 73  
PI WO 9747192 A1 19971218 (199807)\* EN 53p 11  
RW: AT BE CH DE DK EA ES FI FR GB GR IE IT KE LS LU MC MW NL OA PT SD  
SE SZ UG  
W: AL AM AT AU AZ BB BG BR BY CA CH CN CZ DE DK EE ES FI GB GE HU IL  
IS JP KE KG KP KR KZ LK LR LS LT LU LV MD MG MK MN MW MX NO NZ PL  
PT RO RU SD SE SG SI SK TJ TM TR TT UA UG UZ VN  
AU 9661412 A 19980107 (199820)  
ZA 9710452 A 19980826 (199840)# 79p  
US 5879875 A 19990309 (199917)#  
NZ 333030 A 20000526 (200033)  
EP 1018866 A1 20000719 (200036) EN  
R: CH DE FR GB LI SE  
JP 2000512625 W 20000926 (200051) 50p  
AU 725247 B 20001012 (200055)  
AU 2001010037 A 20010315 (200121)#  
AU 742402 B 20020103 (200209)#  
EP 1018866 B1 20021204 (200303) EN  
R: CH DE FR GB LI SE  
DE 69625254 E 20030116 (200313)  
ADT WO 9747192 A1 WO 1996-NZ57 19960614; AU 9661412 A AU 1996-61412 19960614,  
WO 1996-NZ57 19960614; ZA 9710452 A ZA 1997-10452 19971120; US 5879875 A  
US 1996-662244 19960614; NZ 333030 A NZ 1996-333030 19960614, WO 1996-NZ57  
19960614; EP 1018866 A1 EP 1996-918938 19960614, WO 1996-NZ57 19960614; JP  
2000512625 W WO 1996-NZ57 19960614, JP 1997-541244 19960614; AU 725247 B  
AU 1996-61412 19960614, WO 1996-NZ57 19960614; AU 2001010037 A Div ex AU  
1996-61412 19960614, AU 2001-10037 20010103; AU 742402 B Div ex AU  
1996-61412 19960614, AU 2001-10037 20010103; EP 1018866 B1 EP 1996-918938  
19960614, WO 1996-NZ57 19960614; DE 69625254 E DE 1996-625254 19960614, EP  
1996-918938 19960614, WO 1996-NZ57 19960614  
FDT AU 9661412 A Based on WO 9747192; NZ 333030 A Based on WO 9747192; EP  
1018866 A1 Based on WO 9747192; JP 2000512625 W Based on WO 9747192; AU  
725247 B Previous Publ. AU 9661412, Based on WO 9747192; AU 2001010037 A  
Div ex AU 725247; AU 742402 B Previous Publ. AU 200110037, Div ex AU  
725247; EP 1018866 B1 Based on WO 9747192; DE 69625254 E Based on EP  
1018866, Based on WO 9747192  
PRAI WO 1996-NZ57 19960614; ZA 1997-10452 19971120; US 1996-662244  
19960614; AU 2001-10037 20010103  
AB WO 9747192 A UPAB: 20030224  
The following are claimed: (1) a **solution** for preservation of  
biological materials: (A) comprising: (a) a first neutral solute with a  
molecular weight of at least 335 and a solubility in water of at least 0.3  
M, and (b) a second neutral solute with a molecular weight < 200 and with  
both hydrophilic and hydrophobic moieties; or (B) which is isotonic with  
the biological materials and is free of univalent oxyanions and iodide,  
and (2) preservation of biological materials by: (a) pretreating the  
biological material with a **solution** which includes sodium  
butyrate, and (b) contacting the biological material with a preservative  
**solution**.

The **solution** (A) is free of univalent oxyanions and iodide,  
and also comprises at least 1 ion from a protein-stabilising end of the  
Hofmeister series. The first solute is selected from disaccharides and  
trisaccharides, especially raffinose, trehalose, sucrose, lactose

and their analogues. The second neutral solute is selected from trimethylamine oxide, betaine, taurine, sarcosine, **glucose**, mannose, fructose, ribose, galactose, sorbitol, **mannitol**, inositol and their analogues. The **solution** may also comprise sodium sulphate. It may also comprise calcium, which is present as calcium sulphate at a concentration of 1.5-2.0 mM. The **solution** is in a concentrated form, especially in the form of a solid. Components (a) and (b) are typically present at a ratio of 1.4-1.8:1.

USE - The **solutions** may be used for preservation of materials such as **organs**, **tissues** and **cells** from mammals, marine organisms and plants. They may be used e.g. in treatment of leukaemia. In this case, bone marrow is removed from a patient and contacted with the **solution** for at least 3 days, in order to purge the bone marrow of leukaemic **cells**. The purged bone marrow is then returned to the patient.

ADVANTAGE - The **solutions** are of low toxicity, resulting in fewer side effects when biological materials, such as transplant **organs**, are returned to a patient.

Dwg.0/20



L69 ANSWER 5 OF 39 WPIDS (C) 2003 THOMSON DERWENT

AN 2002-479639 [51] WPIDS

DNC C2002-136479

TI Vitrifaction of natural or engineered **tissue** or **organ** other than blood vessel involves immersing **tissue** in cryoprotectant **solutions** with increasing concentrations and cooling to below glass transition temperature.

DC A96 D22 E19 G04 J07

IN BROCKBANK, K G M; KHIRABADI, B S; SONG, Y C

PA (ORGA-N) ORGAN RECOVERY SYSTEMS

CYC 94

PI WO 2002032225 A2 20020425 (200251)\* EN 32p

RW: AT BE CH CY DE DK EA ES FI FR GB GH GM GR IE IT KE LS LU MC MW MZ

NL OA PT SD SE SL SZ TR TZ UG ZW

W: AE AG AL AM AT AU AZ BA BB BG BR BY BZ CA CH CN CO CR CU CZ DE DK

DM DZ EE ES FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ

LC LK LR LS LT LU LV MA MD MG MK MN MW MX MZ NO NZ PL PT RO RU SD

SE SG SI SK SL TJ TM TR TT TZ UA UG UZ VN YU ZA ZW

AU 2002011792 A 20020429 (200255)

ADT WO 2002032225 A2 WO 2001-US32415 20011018; AU 2002011792 A AU 2002-11792 20011018

FDT AU 2002011792 A Based on WO 200232225

PRAI US 2000-691197 2000/10/19

AB WO 200232225 A UPAB: 20020812

NOVELTY - A natural or engineered **tissue** or **organ** other than a blood vessel is vitrified by:

(i) immersing it in a series of **solutions** having increasing concentrations of cryoprotectant to achieve a cryoprotectant concentration for vitrification;

(ii) rapidly cooling to between -80 deg. C and the glass transition temperature (T<sub>g</sub>); and

(iii) further cooling to below T<sub>g</sub>.

DETAILED DESCRIPTION - Vitrifaction of a natural or engineered **tissue** or **organ** other than a blood vessel comprises:

(i) immersing the **tissue** (3) or **organ** in a series of **solutions** having increasing concentrations of cryoprotectant and each having a temperature above -15 deg. C;

(ii) cooling the **tissue** of **organ** at 2.5-100 deg. C per minute from a temperature above -15 deg. C to between -80 deg. C and the T<sub>g</sub>; and

(iii) further cooling at average rate less than 30 deg. C per minute from between -80 deg. C and the T<sub>g</sub> to below the T<sub>g</sub> to vitrify the **tissue** or **organ**.

An INDEPENDENT CLAIM is also included for a method for removing a **tissue** or **organ** other than a blood from vitrification in a **solution** containing cryoprotectant by:

(a) warming the vitrified **tissue** or **organ** in a **solution** containing cryoprotectant at an average rate of 20-40 deg. C per minute to between -80 deg. C and the T<sub>g</sub>;

(b) further warming in the **solution** at an average rate of 200-300 deg. C per minute to a temperature above -75 deg. C; and

(c) immersing the **tissue** or **organ** in a series of **solutions** having decreasing concentrations of cryoprotectant.

USE - For vitrifying a natural or engineered **tissue** or **organ** other than a blood vessel, particularly musculoskeletal **tissue**, cartilage, menisci, muscles, ligaments, tendons, skin, cardiovascular **tissue**, heart valves, myocardium, periodontal **tissue**, glandular **tissue**, islets of Langerhans, cornea, ureter, urethra, pancreas, bladder, kidney, breast, liver, intestine or heart.

ADVANTAGE - The vitrification method results in a greater number or percentage of viable **cells** in a **tissue** or **organ** sample, compared to conventional cryopreservation

techniques. It results in **tissue** or **organ** samples having at least 50% viable **cells**.

DESCRIPTION OF DRAWING(S) - The figure shows a perfusion system that can be used in the invention.

**Tissue 3**

Dwg.1/4

TI     Vitrification of natural or engineered **tissue** or **organ** other than blood vessel involves immersing **tissue** in cryoprotectant **solutions** with increasing concentrations and cooling to below glass transition temperature.

AB     WO 200232225     UPAB: 20020812

NOVELTY - A natural or engineered **tissue** or **organ** other than a blood vessel is vitrified by:

(i) immersing it in a series of **solutions** having increasing concentrations of cryoprotectant to achieve a cryoprotectant concentration for vitrification;

(ii) rapidly cooling to between -80 deg. C. . . glass transition temperature (Tg); and

(iii) further cooling to below Tg.

DETAILED DESCRIPTION - Vitrification of a natural or engineered **tissue** or **organ** other than a blood vessel comprises:

(i) immersing the **tissue** (3) or **organ** in a series of **solutions** having increasing concentrations of cryoprotectant and each having a temperature above -15 deg. C;

(ii) cooling the **tissue** of **organ** at 2.5-100 deg. C per minute from a temperature above -15 deg. C to between -80 deg. C and the . . . 30 deg. C per minute from between -80 deg. C and the Tg to below the Tg to vitrify the **tissue** or **organ**.

An INDEPENDENT CLAIM is also included for a method for removing a **tissue** or **organ** other than a blood from vitrification in a **solution** containing cryoprotectant by:

(a) warming the vitrified **tissue** or **organ** in a **solution** containing cryoprotectant at an average rate of 20-40 deg. C per minute to between -80 deg. C and the Tg;

(b) further warming in the **solution** at an average rate of 200-300 deg. C per minute to a temperature above -75 deg. C; and

(c) immersing the **tissue** or **organ** in a series of **solutions** having decreasing concentrations of cryoprotectant.

USE - For vitrifying a natural or engineered **tissue** or **organ** other than a blood vessel, particularly musculoskeletal **tissue**, cartilage, menisci, muscles, ligaments, tendons, skin, cardiovascular **tissue**, heart valves, myocardium, periodontal **tissue**, glandular **tissue**, islets of Langerhans, cornea, ureter, urethra, pancreas, bladder, kidney, breast, liver, intestine or heart.

ADVANTAGE - The vitrification method results in a greater number or percentage of viable **cells** in a **tissue** or **organ** sample, compared to conventional cryopreservation techniques. It results in **tissue** or **organ** samples having at least 50% viable **cells**.

DESCRIPTION OF DRAWING(S) - The figure shows a perfusion system that can be used in the invention.

**Tissue 3**

Dwg.1/4

TECH

UPTX: 20020812

TECHNOLOGY FOCUS - BIOTECHNOLOGY - Preferred Method: Cooling the **tissue** or **organ** from above -15 degreesC to between -80 degreesC and the Tg is performed at an average rate of at least 10 degreesC (preferably 30-60 degreesC) per minute. Cooling the **tissue** or **organ** from between -80 degreesC and the Tg to below the Tg is performed at an average rate less than 10 degreesC per minute. The immersion step (i) comprises immersing the **tissue** or **organ** in a cryoprotectant-free **solution**; immersing the **tissue** or **organ** in **solution(s)** containing

cryoprotectant at a concentration less than the concentration sufficient for the vitrification; and immersing the **tissue** or **organ** in a **solution** containing cryoprotectant at the concentration sufficient for vitrification. In each immersion step, the **tissue** or **organ** is immersed in the **solution** for at least 10 minutes. The increasing and decreasing concentrations of cryoprotectant are (a) 5-50%, 15-35%, 40-60%, and 65-85%; or (b) 40-60%, 30-45%, 15-35%, 5-20%, 2.5-10%, and 0%.

TECHNOLOGY FOCUS - ORGANIC CHEMISTRY - Preferred Components: The cryoprotectant **solution** comprises acetamide, agarose, alginate, alanine, albumin, ammonium acetate, anti-freeze proteins, butanediol, chondroitin sulfate, chloroform, choline, cyclohexanediols, dextrans, diethylene glycol, dimethyl acetamide, dimethyl formamide, dimethyl sulfoxide, erythritol, ethanol, ethylene glycol, ethylene glycol monomethyl ether, formamide, **glucose**, glycerol, glycerophosphate, glyceryl monoacetate, glycine, glycoproteins, hydroxyethyl starch, inositol, **lactose**, magnesium chloride, magnesium sulfate, maltose, **mannitol**, mannose, methanol, methoxy propanediol, methyl acetamide, methyl formamide, methyl ureas, methyl **glucose**, methyl glycerol, phenol, pluronic polyols, polyethylene glycol, polyvinyl pyrrolidone, proline, 1,2-propanediol, pyridine N-oxide, raffinose, ribose, serine, sodium bromide, sodium chloride, . . . and/or xylose. It preferably comprises (in weight per volume) dimethyl sulfoxide (20-30%), formamide (10-20%) and 1,2-propanediol (10-20%) in a vehicle **solution**. Each **solution** comprises an osmotic buffering agent, preferably **mannitol**.

TT TT: VITREOUS NATURAL ENGINEERING **TISSUE ORGAN BLOOD**  
VESSEL IMMERSE **TISSUE SOLUTION** INCREASE  
CONCENTRATE COOLING BELOW GLASS TRANSITION TEMPERATURE.

=>

L7 ANSWER 23 OF 37 CA COPYRIGHT 2003 ACS  
AN 121:252473 CA  
TI Effect of polyols on the post-thawing motility of pellet-frozen ram spermatozoa  
AU Molinia, F. C.; Evans, G.; Maxwell, W. M. C.  
CS Department Animal Science, University Sydney, Sydney, NSW 2006, Australia  
SO Theriogenology (1994), 42(1), 15-23  
CODEN: THGNBO; ISSN: 0093-691X  
DT Journal  
LA English  
AB The cryoprotective effects of polyols in the absence and presence of glycerol in Tris-glucose-egg yolk based diluents on the post-thawing motility and acrosome integrity of pellet-frozen ram spermatozoa were examd. Incorporation of adonitol or xylitol (low mol. wt. polyols; LMWPs) in diluents improved motility of spermatozoa in the absence of glycerol with max. motility at 0.3 M (26.9 vs. 13.3% mean motile spermatozoa). Five concns. of adonitol (0, 150, 300, 450, 600 mM) were examd. in diluents contg. 5 concns. of glycerol (0, 1.5, 3.0, 4.5, 6.0% vol./vol.). There was an additive effect of incorporation of 1.5% vol./vol. glycerol with up to 450 mM adonitol, but at higher levels of glycerol the incorporation of adonitol was detrimental to motility. The acrosome integrity of spermatozoa in diluents contg. 0, 150 and 300 mM adonitol was superior to those contg. 450 and 600 mM adonitol (46.1 vs. 35.1% mean intact acrosomes). Among the high mol. wt. polyols (HMWPs) examd., better recovery of spermatozoa was obtained in diluents contg. sorbitol than mannitol or inositol. Sorbitol or mannitol (300 mM) improved the motility of spermatozoa in diluents without glycerol, but the incorporation of HMWPs was detrimental in diluents contg. glycerol. All five polyols were examd. in isotonic diluents contg. 360:0, 300:55, 240:110, 180:165, 120:220mM (Tris:polyol; 360 mosmol) and 6.0% vol./vol. glycerol. There was a linear decrease in motility and acrosome integrity of spermatozoa with increasing polyol concn. in the diluent except for inositol, which was not detrimental. The authors conclude that the polyols examd. have a cryoprotective effect on pellet-frozen ram spermatozoa except for inositol. However, in the authors study, no combination of polyols and glycerol was superior in terms of post-thawing motility and acrosome integrity of spermatozoa to 6.0% vol./vol. glycerol alone in Tris-glucose-egg yolk diluents.

L1 ANSWER 1 OF 1 REGISTRY COPYRIGHT 2002 ACS  
RN 9041-07-0 REGISTRY  
CN **Decaglycerol (9CI)** (CA INDEX NAME)  
OTHER NAMES:  
CN Decaglycerin  
CN Polyglycerin 10  
DR 83689-42-3, 26085-10-9, 34322-27-5  
MF C30 H62 O21  
CI IDS, COM, MAN  
LC STN Files: BIOBUSINESS, BIOSIS, CA, CAPLUS, CASREACT, CHEMLIST, IFICDB,  
IFIPAT, IFIUDB, NIOSHTIC, TOXCENTER, USPATFULL  
Other Sources: EINECS\*\*, NDSL\*\*, TSCA\*\*  
(\*\*Enter CHEMLIST File for up-to-date regulatory information)

\*\*\* STRUCTURE DIAGRAM IS NOT AVAILABLE \*\*\*  
127 REFERENCES IN FILE CA (1962 TO DATE)  
63 REFERENCES TO NON-SPECIFIC DERIVATIVES IN FILE CA  
127 REFERENCES IN FILE CAPLUS (1962 TO DATE)

=>

L9 ANSWER 114 OF 268 WPIDS (C) 2002 THOMSON DERWENT

AN 1998-076784 [07] WPIDS

CR 1998-593980 [50]

DNC C1998-025629

TI **Solution** for preservation of biological materials - comprises two neutral solutes, especially raffinose and tri methylamine oxide.

DC B04 D16 D22 E19 E37 P31

IN FERGUSON, A B; WIGGINS, P M; WATSON, J D

PA (BIOS-N) BIOSTORE NEW ZEALAND LTD; (BIOS-N) BIOSTORE NZ LTD; (BIOS-N) BIOSTORE NEW ZEALAND

CYC 73

PI WO 9747192 A1 19971218 (199807)\* EN 53p A01N001-00

RW: AT BE CH DE DK EA ES FI FR GB GR IE IT KE LS LU MC MW NL OA PT SD  
SE SZ UG

W: AL AM AT AU AZ BB BG BR BY CA CH CN CZ DE DK EE ES FI GB GE HU IL  
IS JP KE KG KP KR KZ LK LR LS LT LU LV MD MG MK MN MW MX NO NZ PL  
PT RO RU SD SE SG SI SK TJ TM TR TT UA UG UZ VN

AU 9661412 A 19980107 (199820) A01N001-00

ZA 9710452 A 19980826 (199840)# 79p A61K000-00

US 5879875 A 19990309 (199917)# A01N001-02

NZ 333030 A 20000526 (200033) A01N003-00

EP 1018866 A1 20000719 (200036) EN A01N001-00

R: CH DE FR GB LI SE

JP 2000512625 W 20000926 (200051) 50p A01N001-00

AU 725247 B 20001012 (200055) A01N001-00

AU 2001010037 A 20010315 (200121)# A61K035-28

AU 742402 B 20020103 (200209)# A61K035-28

ADT WO 9747192 A1 WO 1996-NZ57 19960614; AU 9661412 A AU 1996-61412 19960614,  
WO 1996-NZ57 19960614; ZA 9710452 A ZA 1997-10452 19971120; US 5879875 A  
US 1996-662244 19960614; NZ 333030 A NZ 1996-333030 19960614, WO  
1996-NZ57

19960614; EP 1018866 A1 EP 1996-918938 19960614, WO 1996-NZ57 19960614;

JP

2000512625 W WO 1996-NZ57 19960614, JP 1997-541244 19960614; AU 725247 B  
AU 1996-61412 19960614, WO 1996-NZ57 19960614; AU 2001010037 A Div ex AU  
1996-61412 19960614, AU 2001-10037 20010103; AU 742402 B Div ex AU  
1996-61412 19960614, AU 2001-10037 20010103

FDT AU 9661412 A Based on WO 9747192; NZ 333030 A Based on WO 9747192; EP  
1018866 A1 Based on WO 9747192; JP 2000512625 W Based on WO 9747192; AU  
725247 B Previous Publ. AU 9661412, Based on WO 9747192; AU 2001010037 A  
Div ex AU 725247; AU 742402 B Previous Publ. AU 200110037, Div ex AU  
725247

PRAI WO 1996-NZ57 19960614; ZA 1997-10452 19971120; US 1996-662244  
19960614; AU 2001-10037 20010103

IC ICM A01N001-00; A01N001-02; A01N003-00; A61K000-00; A61K035-28

ICS A61B000-00; A61K031-00; A61K035-12; A61K035-14; A61K035-34;  
A61K035-36; A61K035-60; A61K035-78; A61P035-02; C12N001-04;  
C12N005-00; C12N005-06

AB WO 9747192 A UPAB: 20020208

The following are claimed: (1) a **solution** for preservation of  
biological materials: (A) comprising: (a) a first neutral solute with a  
molecular weight of at least 335 and a solubility in water of at least

0.3

M, and (b) a second neutral solute with a molecular weight < 200 and with  
both hydrophilic and hydrophobic moieties; or (B) which is isotonic with  
the biological materials and is free of univalent oxyanions and iodide,  
and (2) preservation of biological materials by: (a) pretreating the  
biological material with a **solution** which includes sodium  
butyrate, and (b) contacting the biological material with a preservative

**solution.**

The **solution** (A) is free of univalent oxyanions and iodide, and also comprises at least 1 ion from a protein-stabilising end of the Hofmeister series. The first solute is selected from disaccharides and trisaccharides, especially raffinose, trehalose, sucrose, **lactose** and their analogues. The second neutral solute is selected from trimethylamine oxide, betaine, taurine, sarcosine, glucose, mannose, fructose, ribose, galactose, sorbitol, **mannitol**, inositol and their analogues. The **solution** may also comprise sodium sulphate. It may also comprise calcium, which is present as calcium sulphate at a concentration of 1.5-2.0 mM. The **solution** is in a concentrated form, especially in the form of a solid. Components (a) and (b) are typically present at a ratio of 1.4-1.8:1.

USE - The **solutions** may be used for preservation of materials such as organs, tissues and cells from mammals, marine organisms

and plants. They may be used e.g. in treatment of leukaemia. In this case, bone marrow is removed from a patient and contacted with the **solution** for at least 3 days, in order to purge the bone marrow of leukaemic cells. The purged bone marrow is then returned to the patient.

ADVANTAGE - The **solutions** are of low toxicity, resulting in fewer side effects when biological materials, such as transplant organs, are returned to a patient.

Dwg.0/20

FS CPI GMPI

FA AB; DCN

MC CPI: B04-C02X; B10-A07; B14-H01A; B14-N17B; D05-H01; D09-A01; E07-A02A; E07-A02D; E07-A02H; E10-A03

L28 ANSWER 30 OF 31 CA COPYRIGHT 2002 ACS  
AN 98:213494 CA  
TI Identification of new cryoprotective agents for cultured mammalian cells  
AU Klebe, Robert J.; Mancuso, Melodee G.  
CS Dep. Anat., Univ. Texas Health Sci. Cent., San Antonio, TX, 78284, USA  
SO In Vitro (1983), 91(3, pt. 1), 167-70  
CODEN: ITCSAF; ISSN: 0073-5655  
DT Journal  
LA English  
AB Thirty-one compds. were identified that act as cryoprotective agents for cultured mammalian (CHO) cells. Eight compds. were comparable to DMSO in cryoprotective effectiveness. Many of the cryoprotective compds. studied also (1) promote cell fusion and (2) induce cell differentiation in erythroleukemia and other cell systems. Thus, previously unrecognized effects on the differentiated state of cells may occur when cells are treated with cryoprotective agents.



ANSWER 7 OF 30 MEDLINE

AN 1998353576 MEDLINE  
DN 98353576 PubMed ID: 9687336  
TI The physical state of **mannitol** after freeze-drying: effects of **mannitol** concentration, freezing rate, and a noncrystallizing cosolute.  
AU Kim A I; Akers M J; Nail S L  
CS Department of Industrial and Physical Pharmacy, Purdue University, West Lafayette, Indiana 47907, USA.  
SO JOURNAL OF PHARMACEUTICAL SCIENCES, (1998 Aug) 87 (8) 931-5.  
Journal code: 2985195R. ISSN: 0022-3549.  
CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 199809  
ED Entered STN: 19980910  
Last Updated on STN: 19980910  
Entered Medline: 19980903  
AB The objectives of this study were to (1) measure the effects of freezing rate and **mannitol** concentration on the physical state of freeze-dried **mannitol** when **mannitol** is present as a single component, (2) determine the relative concentration threshold above which crystalline **mannitol** can be observed by X-ray powder diffraction in the freeze-dried solid when a variety of noncrystallizing solutes are included in the formulation, and (3) measure the glass transition temperature of amorphous **mannitol** and to determine the degree to which the glass transition temperature of freeze-dried solids consisting of **mannitol** and a disaccharide is predicted by the Gordon-Taylor equation. Both freezing rate and **mannitol** concentration influence the crystal form of **mannitol** in the freeze-dried solid when **mannitol** is present as a single component. Slow freezing of 10% (w/v) **mannitol** produces a mixture of the alpha and beta polymorphs, whereas fast freezing of the same solution produces the delta form. Fast freezing of 5% (w/v) **mannitol** results primarily in the beta form. The threshold concentration above which crystalline **mannitol** is detected in the freeze-dried solid by X-ray diffraction is consistently about 30% (w/w) when a second, noncrystallizing solute is present, regardless of the nature of the second component. The glass transition temperature of amorphous **mannitol** measured from the quench-cooled melt is approximately 13 degreesC. Accordingly, **mannitol** is an effective plasticizer of freeze-dried solids when the **mannitol** remains amorphous. Glass transition temperatures of mixtures of **mannitol** and the disaccharides sucrose, maltose, trehalose, and **lactose** are well predicted by the Gordon-Taylor equation with values of k in the range of 3 to 4.

~~9041-07-0~~  
~~9002-89-5~~ \*

30, 28, 25, 13, 10, 8

L  
L9 ANSWER 184 OF 268 WPIDS (C) 2002 THOMSON DERWENT

AN 1989-162877 [22] WPIDS

DNC C1989-072372

TI New blood-preservative liq. compsn. - contg. haemolysis preventive of mono

carboxylic ester and other blood-preservative liq. ingredients, esp. for red blood corpuscles.

DC B05 D22 E19

PA (KAOS) KAO CORP; (TERU) TERUMO CORP

CYC 1

PI JP 01106826 A 19890424 (198922)\* 8p

ADT JP 01106826 A JP 1987-263695 19871021

PRAI JP 1987-263695 19871021

IC A61K035-14

AB JP 01106826 A UPAB: 19930923

New blood-preservative liq. compsn. contains a haemolysis preventive of monocarboxylic ester of formula  $\text{RCOOR}'$  (I) and other blood-preservative liq. ingredients. (R and R' (individually) are 3C or higher chain hydrocarbon with  $\text{R} + \text{R}' = 11-30$ ). Blood-preservative liqs. are e.g. anticoagulants and at least one of sodium citrate, citric acid, glucose, monosodium phosphate, adenine, sodium chloride, mannitol, maltose, multitol, sorbitol, sucrose, and lactose. Compsn. is prepd. by blending (1) with base liq. of ACD, CPD, CPDA-1, CPDA-2, SAG

and

SAG solns. contg. mannitol, maltose, maltose, multitol, sorbitol, sucrose, and lactose. Either or both of R and R' are pref. branched. Final concn. of ester is 100 M - 10mM; and concn. is 400microM - 4 mM for branched R and/or R'.

USE/ADVANTAGE - Compsn. with high physiological safety, has good protective action esp. on red blood corpuscles. Effect is durable.  
0/0

FS